

Maryville Christian School
Preseason Parental Information and Consent Form for Concussions

What is a concussion?

A concussion is an injury to the brain. It is caused by a bump, blow, or jolt to either the head or the body that causes the brain to move rapidly within the skull. The resulting injury to the brain changes how the brain functions in a normal manner. The signs and symptoms of a concussion can show up immediately after the injury or may not appear for hours or days after the injury. Concussions can have serious long-term health effects, and even a seemingly mild injury can be serious. A major concern with any concussion is returning to play too soon. Having a second concussion before healing can take place from the initial or previous concussion can lead to serious and potentially fatal health conditions.

What are the symptoms of a concussion?

Signs and symptoms of a concussion are typically noticed right after the injury, but some might not be recognized until days after the injury. Common symptoms include: headache, dizziness, amnesia, fatigue, confusion, mood changes, depression, poor vision, sensitivity to light or noise, lethargy, poor attention or concentration, sleep disturbances, and aggression. The individual may or may not have lost consciousness.

What should the athlete know about playing with a concussion?

Teach athletes that it's not smart to play with a concussion. Rest is the key after a concussion. Sometimes athletes, parents, and other school or league officials wrongly believe that it shows strength and courage to play injured. Discourage others from pressuring injured athletes to play. Don't let your athlete convince you that they are "just fine."

What are the risks of returning to activity too soon after sustaining a concussion?

Returning too soon can create long-term problems. If an athlete has a concussion, their brain needs time to heal. Don't let him or her return to play the day of the injury and until a health care professional, experienced in evaluating for concussion, says he or she is symptom-free and it's OK to return to play. A repeat concussion that occurs before the brain recovers from the first—usually within a short time period (hours, days, weeks)—can slow recovery or increase the chances for long-term problems.

What can happen if my child keeps on playing with a concussion?

Athletes with the signs and symptoms of a concussion should be removed from play immediately. Continuing to play with the signs and symptoms of a concussion leaves the young athlete especially vulnerable to greater injury. There is an increased risk of significant damage from a concussion for a period of time after that concussion occurs, particularly if the athlete suffers another concussion before completely recovering from the first one. This can lead to prolonged recovery, or even to severe brain swelling (second impact syndrome) with devastating and even fatal consequences. It is well known that adolescent or teenage athletes will often under report symptoms of injuries, and concussions are no different. As a result, education of administrators, coaches, parents and students is the key for the student-athlete's safety.

Parental Consent

By signing this form, I understand the risks and dangers related with returning to play too soon after a concussion. Furthermore, in the event that my son/daughter is diagnosed with a concussion, I give my consent for my son/daughter to participate in and comply with the Maryville Christian School return to play protocol. The undersigned, being a parent, guardian, or another person with legal authority, grants this permission. If I refuse to consent to the MCS protocol, my child WILL NOT be allowed to participate in Maryville Christian School Athletics.

Athlete's Name (print)

Parent's or Guardian's Name (print)

Parent's or Guardian's Signature

Date: _____